

CHAPTER 3

Genetically encoded molecular tools for light-driven silencing of targeted neurons

Brian Y. Chow[†], Xue Han[‡] and Edward S. Boyden^{§,¶,||,*}

[†] Department of Bioengineering, University of Pennsylvania, Philadelphia, PA, USA

[‡] Department of Biomedical Engineering, Boston University, Boston, MA, USA

[§] MIT Media Lab, MIT, Cambridge, MA, USA

[¶] Department of Biological Engineering, MIT, Cambridge, MA, USA

^{||} Department of Brain and Cognitive Sciences, MIT, Cambridge, MA, USA

Abstract: The ability to silence, in a temporally precise fashion, the electrical activity of specific neurons embedded within intact brain tissue, is important for understanding the role that those neurons play in behaviors, brain disorders, and neural computations. “Optogenetic” silencers, genetically encoded molecules that, when expressed in targeted cells within neural networks, enable their electrical activity to be quieted in response to pulses of light, are enabling these kinds of causal circuit analyses studies. Two major classes of optogenetic silencer are in broad use in species ranging from worm to monkey: light-driven inward chloride pumps, or halorhodopsins, and light-driven outward proton pumps, such as archaerhodopsins and fungal light-driven proton pumps. Both classes of molecule, when expressed in neurons via viral or other transgenic means, enable the targeted neurons to be hyperpolarized by light. We here review the current status of these sets of molecules, and discuss how they are being discovered and engineered. We also discuss their expression properties, ionic properties, spectral characteristics, and kinetics. Such tools may not only find many uses in the quieting of electrical activity for basic science studies but may also, in the future, find clinical uses for their ability to safely and transiently shut down cellular electrical activity in a precise fashion.

Keywords: optogenetics; opsins; neural silencing; halorhodopsin; archaerhodopsin; channelrhodopsin; control; cell types; neural circuits; causality.

Introduction

The ability to silence, in a temporally precise fashion, the electrical activity of specific neurons embedded within intact brain tissue, is important

*Corresponding author.
Tel.: +617 324-3085
E-mail: eboyden3@gmail.com

for understanding the role that those neurons play in behaviors, brain disorders, and neural computations. Over the past several years, we and others have discovered that a large number of light-driven ion pumps naturally occurring in archaea, fungi, and other species, can be genetically expressed in neurons, enabling them to be hyperpolarized in response to light. These molecules, microbial (type I) opsins, are seven-transmembrane proteins, which translocate specific ions from one side of the membrane in which they are expressed, to the other, in response to light. For example, halorhodopsins are light-driven inward chloride pumps, which, when expressed in neurons, enable them to be hyperpolarized by orange light (Fig. 1ai). Archaeorhodopsins are light-driven outward proton pumps which, when expressed in neurons, also result in hyperpolarization (Fig. 1bi). These proteins are monolithic, and encoded for by relatively small genes, under a kilobase long, small enough that genetic delivery to specific cells (e.g., via viruses, transfection, electroporation, transgenesis, or other methods) is achievable using many conventional methods in animals commonly used in neuroscience, from *Caenorhabditis elegans* to nonhuman primate (Adamantidis et al., 2007; Atasoy et al., 2008; Bi et al., 2006; Chan et al., 2010; Chhatwal et al., 2007; Dittgen et al., 2004; Douglass et al., 2008; Han et al., 2009a,b; Huber et al., 2008; Ishizuka et al., 2006; Kuhlman and Huang, 2008; Lagali et al., 2008; Li et al., 2005; Nagel et al., 2005; Petreanu et al., 2007; Schroll et al., 2006; Tan et al., 2008; Toni et al., 2008; Wang et al., 2007; Zhang et al., 2006). Further, thanks to the proliferation of optical devices driven by microscopy innovation as well as from other fields (e.g., telecommunications and computing), many devices exist or are easily engineered for the delivery of light to specific brain regions, even deep within the mammalian brain (conveyed via implanted optical fibers or waveguides; e.g., a typical 200- μm diameter fiber with end-of-tip irradiance of 100–200mW/mm² can illuminate a volume of a few cubic millimeters), and even in a wirelessly

controlled fashion (Aravanis et al., 2007; Gradinaru et al., 2007; Wentz et al., 2011; Zorzos et al., 2010). *In vitro*, conventional microscope fluorescence illuminators, such as xenon lamps, mercury lamps, confocal or two-photon lasers, and LEDs, equipped with fast shutters, deflectors, or controllers (Boyden et al., 2005; Campagnola et al., 2008; Farah et al., 2007; Guo et al., 2009; Petreanu et al., 2007, 2009; Rickgauer and Tank, 2008), suffice to drive these molecules, the majority of which operate under illumination with incident light in the range of 0.1–10mW/mm² at typical levels of expression in typical cells.

The speed of operation of these molecules is high, as they respond to light within milliseconds, and shut off rapidly after cessation of light delivery. For operation, these proteins require the cofactor all-*trans*-retinal as their essential chromophore, which acts to capture light. This chemical, being a natural beta carotene derivative, occurs in many organisms (such as mice, rats, and primates) at high enough background levels so that external chemical supplementation is not needed; for species such as *C. elegans* and *Drosophila*, whose background levels are lower, feeding of the all-*trans*-retinal to the organism suffices to enable supplementation. Because of this ease of use, adoption of these molecules for use in neuroscience has been rapid.


Overview of light-driven ion pumps

The first fully genetically encoded optogenetic tool to be utilized in neuroscience was the light-gated cation channel channelrhodopsin-2 (ChR2) from the green alga *Chlamydomonas reinhardtii*. This light-gated nonspecific cation channel, when expressed in neurons, enable the neurons to be depolarized with blue light (Boyden et al., 2005; Nagel et al., 2003). The channelrhodopsins, and the origins of optogenetic tools, are reviewed elsewhere in this volume (Chapter 2), as well as in other earlier references (Boyden, 2011; Chow et al., 2011a). Like halorhodopsins and light-driven

(a)


i

Halorhodopsins
(e.g., Halo / NpHR)



ii

Halorhodopsin-mediated
silencing of cultured hippocampal
neuron spiking activity



iii


Orexin/hypocretin neuron
expressing Halo, and
silenced by yellow light



(b)


i

Archaeorhodopsins
(e.g., Arch, ArchT)




ii

Arch photocurrents




ArchT photocurrents



iii

Cortical pyramidal neuron
expressing Arch, and
silenced by yellow light



proton pumps, channelrhodopsins are all-*trans*-retinal binding seven-transmembrane proteins, and so there are some similarities in structure and function across these different opsin classes. Unlike channelrhodopsins, which open up a channel pore when gated by light, halorhodopsins and light-driven proton pumps translocate one ion per photon absorbed, which may in principle limit light sensitivity, but also means that they can pump ions against a concentration gradient, causing significant voltage swings less limited by the reversal potential of specific ions in neurons.

Halorhodopsins, as exemplified by the light-driven inward chloride pump halorhodopsin from the species *Natronomonas pharaonis* (Halo/NpHR; Lanyi et al., 1990), were found in 2007 to function in mammalian neurons, where they mediate hyperpolarizing currents of 40–100 pA (Han and Boyden, 2007; Zhang et al., 2007), which is sufficient to support modest neural silencing (Fig. 1aii). Because of the low currents that the original Halo could produce in mammalian neurons, due to limited protein trafficking in mammalian neurons that hampers expression at high levels (Chow et al., 2010; Gradinaru et al., 2008; Zhao et al., 2008), usage of the original Halo has largely been confined to use in invertebrates such as *C. elegans*, where it appears to function sufficiently in neurons and other excitable cells, to modulate behavior (Zhang et al.,

2007). In 2011, however, a group reported for the first time the use of the original Halo to mediate behavioral changes in living mammals (Tsunematsu et al., 2011), expressing Halo in the orexin/hypocretin neurons of the mouse brain via the generation of a transgenic mouse line. The orexin/hypocretin neurons of this mouse, when illuminated, undergo reductions in spiking (Fig. 1aiii), and when such illumination is performed in the awake mouse, the animals enter slow-wave sleep, thus showing that brief silencing of this pathway is sufficient to induce sleep. Light-driven proton pumps, as exemplified by the archaeal protein archaerhodopsin-3 (Arch/aR-3) from *Halorubrum sodomense* (Fig. 1bi), were found in 2010 to function well in mammalian neurons in the service of light-driven neural silencing (Chow et al., 2010). Arch, and its close relative ArchT, from *Halorubrum* sp. *TP009* (Han et al., 2011), can mediate photocurrents of 900 pA in cultured neurons (Fig. 1bii), and can mediate complete light-driven neural silencing of neural activity in mice and primates (Chow et al., 2010; Han et al., 2011), in response to yellow or green light (Fig. 1biii).

Light-driven inward chloride pumps and light-driven outward proton pumps are found in species of archaea (Ihara et al., 1999; Klare et al., 2008; Lanyi, 2004, 1986; Mukohata et al., 1999), bacteria (Antón et al., 2005; Balashov et al., 2005;

Fig. 1. Two major classes of microbial opsin for light-driven hyperpolarization of neurons. (a) Halorhodopsins, light-driven inward chloride pumps. (i) Diagram of the physiological response of halorhodopsins when expressed in the plasma membranes of neurons and exposed to light. (ii) Light-driven spike blockade, demonstrated for a representative cultured hippocampal neuron (*top*), as well as a population of cultured hippocampal neurons (*bottom*). Neurons expressed Halo/NpHR, and received 20 pulses of somatic current injection (~300 pA, 4 ms, 5 Hz), accompanied by two periods of yellow light delivery (yellow bars). Adapted from Han and Boyden (2007). (iii) A neuron in a transgenic mouse expressing Halo in orexin/hypocretin neurons, being whole-cell patch clamp recorded during yellow light delivery, and exhibiting neural silencing. Adapted from Tsunematsu et al. (2011). (b) Archaerhodopsins, light-driven outward proton pumps. (i) Diagram of the physiological response of archaerhodopsins when expressed in the plasma membranes of neurons and exposed to light. (ii) Photocurrents of Arch (*top*) and ArchT (*bottom*), measured as a function of 575 ± 25 nm light irradiance, in patch-clamped cultured neurons, for low (*left*) and high (*right*) light powers. Adapted from Chow et al. (2010) and Han et al. (2011). (iii) Neural activity in a representative neuron recorded in awake mouse brain (*top, middle*), as well as a population of neurons in awake mouse brain (*bottom*). Shown is neural activity before, during, and after 5 s of yellow light illumination, displayed both as a spike raster plot (*top*) and as a histogram of instantaneous firing rate averaged across trials (*middle, bottom*; bin size, 20 ms). Adapted from Chow et al. (2010).

Beja et al., 2001, 2000; Friedrich et al., 2002; Kelemen et al., 2003; Kim et al., 2008), fungi (Brown, 2004; Waschuk et al., 2005), and algae (Tsunoda et al., 2006). Because halorhodopsins, bacteriorhodopsins, and archaeorhodopsins have been crystallized and characterized by spectroscopy, mutagenesis, and physiology, much is known about their structure–function relationships (Enami et al., 2006; Essen, 2002; Kolbe et al., 2000; Luecke et al., 1999; Yoshimura and Kouyama, 2008). To date, these light-driven ion pumps have been found chiefly by analyzing genomes for sequences that resemble those of light-activated ion pumps and channels, followed by assessment of these gene products in heterologous expression systems (chiefly cell lines, neurons, and intact mouse brain) (Chow et al., 2011b). Interestingly, a great many light-activated ion pumps from species all over the tree of life—even the very first one ever characterized, the *Halobium salinarum* bacteriorhodopsin—were able to mediate neural hyperpolarizations when assessed in cultured mouse neurons (Chow et al., 2010).

Properties of opsins and how they affect performance

Each of these opsins has a different characteristic action spectrum, which describes the set of colors that optimally drive the opsin to function. Although chloride pumps are chiefly driven by yellow light and possess largely stereotyped action spectra, proton pumps exist in many different kingdoms of life that are driven by many different colors of light. For example, the fungal light-driven proton pump from *Leptosphaeria maculans* (Mac) is drivable by blue-green light, and thus, alongside the earlier molecule Halo, which can be driven by yellow-red light, enables two-color silencing of different sets of neurons (Fig. 2b; Chow et al., 2010). That is, a Mac-expressing neuron will be quieted by blue light, but not red light; in contrast, a Halo-expressing neuron will be quieted by red light,




Fig. 2. Silencing of neurons with specific colors of light. (a) Hyperpolarization and depolarization events induced in a representative cultured hippocampal neuron expressing both ChR2 and Halo, by a Poisson train (mean interpulse interval $\lambda=100\text{ms}$) of alternating pulses of yellow and blue light (10 ms duration), denoted by yellow and blue bars, respectively. Adapted from Han and Boyden (2007). (b) Multicolor silencing of two neural populations, enabled by blue- and red-light drivable ion pumps of different genomic classes. Action potentials evoked by current injection into patch-clamped cultured neurons transfected with the archaeal opsin Halo (i) were selectively silenced by the red light but not by blue light, and vice versa in neurons expressing the fungal opsin Mac (ii). Gray boxes in the inset (iii) indicate periods of patch clamp current injection. Adapted from Chow et al. (2010).

and not blue light. ChR2, which is blue light driven, can also be used alongside silencers such as Halo which are yellow or red light driven, even in the same cell, enabling depolarization and hyperpolarization of the doubly targeted cell by blue and yellow light, respectively (Fig. 2a). The delivery of multiple opsins to the same cell can be facilitated by the use of 2A linker peptides to combine the genes for multiple opsins into a

single open reading frame (Han and Boyden, 2007; Han et al., 2009b), useful for bidirectionally assessing the causal role that a given set of neurons plays in a single animal, or for perturbing complex properties of neural dynamics such as neural synchrony (Han and Boyden, 2007).

In this review, we will summarize the properties of these opsins, surveying how they are expressed in cells, how they conduct ions across membranes, how they cycle in response to light, and how they respond to light of different colors and powers. The optogenetic molecules here described were isolated from archaea and fungi, species whose lipid and ionic compositions, and external environment, differ greatly from those of neurons. Accordingly, the heterologous expression of these molecules in neurons may result in performance characteristics that are very different from what would be expected from their expression in their native cells (Chow et al., 2011b). Indeed, this means that the performance of opsins must be assessed directly in the cell types of interest, as it can be difficult to predict how these high-speed molecules will function in a specific milieu, from their properties as characterized in another one.

Some general guidelines and considerations apply, however. First, to insure efficient translation, it is useful to codon-optimize molecules for the target species. Second, to boost membrane expression, it is often helpful to append extra protein sequences to opsins to improve their protein folding, membrane trafficking and localization, and even subcellular compartmentalization (Chow et al., 2010; Gradinaru et al., 2008; Greenberg et al., 2011; Zhao et al., 2008). Third, although many groups use these molecules to change the voltage of neurons, it is also useful to explicitly consider the ions translocated by these pumps; for example, they can be used to change the levels of specific ions in cells or subcellular compartments, in a temporally precise way. Fourth, the photocycle kinetics can be used to determine which molecule might be best for a given scientific application; for example, various light-driven ion pumps may enter different states

that may render them nonfunctional for periods of time (Fig. 4). It is critical to remember that these light-driven ion pumps are not simple on–off switches; they undergo a series of structural rearrangements, with many intermediate states between light reception and final restoration of the molecule to the initial state. Over the past several decades, a myriad of structure–function studies have been performed on these molecules (Bamberg et al., 1993; Blanck and Oesterhelt, 1987; Braiman et al., 1987; Brown et al., 1996; Essen, 2002; Gilles-Gonzalez et al., 1991; Hegemann et al., 1985; Henderson and Schertler, 1990; Kolbe et al., 2000; Lanyi, 2004, 1986; Lanyi et al., 1990; Luecke et al., 1999; Marinetti et al., 1989; Marti et al., 1991; Mogi et al., 1987, 1988, 1989a,b; Rudiger and Oesterhelt, 1997; Subramaniam et al., 1992; Tittor et al., 1997, 1995; Varo et al., 1995a,b), linking specific amino acids with specific kinetic and spectral properties of the molecules. Fifth, the light-sensitivity of a molecule can determine how well it will function in the brain; improving light sensitivity can enable larger volumes of brain to be controlled than possible with less light-sensitive molecules (e.g., Fig. 1bii, *bottom*). In the following sections, we explore these properties, for light-driven chloride pumps and light-driven proton pumps.

Protein expression and photocurrent magnitude of light-driven ion pumps

The *N. pharaonis* halorhodopsin Halo/NpHR, the first microbial opsin to be used for optical neural silencing (Fig. 1a), has a reversal potential of approximately -400mV (Seki et al., 2007), and although it comes from an archaeon that lives in very high salinity environments, the optimal chloride concentration for its operation is actually very close to that found in neurons (in contrast to that of other halorhodopsins such as the *H. salinarum* halorhodopsin, which demands far higher chloride concentrations) (Okuno et al., 1999). This may be one of the reasons why it works in some neurons in some species (Fig. 1a ii and a iii). However, at

high expression levels, Halo forms intracellular aggregates (Gradinaru et al., 2008; Zhao et al., 2008), an issue that can be ameliorated by the appending of trafficking sequences from the Kir2.1 channel (Gradinaru et al., 2008, 2010; Zhao et al., 2008), which reduces aggregation and increases photocurrent manifold over that of the baseline molecule. Other signal sequences, such as a prolactin localization sequence, also increase halorhodopsin photocurrent (Chow et al., 2011a); it is important to note, however, that different sequences may do different things in different cell types in different species.

The *H. sodomense* archaerhodopsin Arch (Fig. 1b), possesses strong photocurrents that exceed the original *N. pharaonis* halorhodopsin currents by an order of magnitude, both at low and at high light power (Fig. 1bii, top) (Chow et al., 2010). Arch can mediate nearly complete silencing of neural activity in awake behaving mice (Fig. 1biii). Arch does not require trafficking sequences to achieve these high levels of performance, although they can help to boost expression and/or current yet further (Chow et al., 2010). Other opsins that are similar in amino acid sequence to Arch have been described in genomic and gene sequence databases; all of these Arch relatives, found in other species of the *Halorubrum* genus, also express well in neurons (Han et al., 2011). One of these Arch relatives, the archaerhodopsin from *Halorubrum* sp. *TP009*, presents photocurrents that are about $3.5\times$ more light sensitive than those of Arch (Fig. 1bii, bottom), supporting the silencing of broad brain regions, and neurons in the cortex of the awake behaving macaque (Han et al., 2011). Light-driven proton pumps are powerful at neural silencing, a perhaps surprising fact given the relative scarcity of protons relative to other major charge carriers in the brain (such as Na^+ , K^+ , or Cl^-); the fast kinetics of archaerhodopsins (Lukashev et al., 1994; Ming et al., 2006), and their good trafficking in mammalian cells, among other attributes, may support the high performance of archaerhodopsins. These light-driven

outward proton pumps, interestingly, do not change cellular pH to a greater extent than do opsins such as ChR2 (whose proton conductance is 10^6 times its sodium conductance). Specifically, illuminating Arch with relatively bright light when expressed in cultured neurons resulted in alkalizations of neurons by 0.1–0.15 pH units, a change in pH of smaller magnitude than that seen with ChR2, perhaps due to the high proton conductance of channelrhodopsins (Berthold et al., 2008; Lin et al., 2009; Nagel et al., 2003). From an end-user standpoint, illumination of Arch- and ArchT-expressing neurons, in awake mice and macaques, for periods of many minutes, did not alter spike waveform or spike frequency when comparing before versus after silencing.

Kinetics of light-driven ion pumps

From a kinetic standpoint, all of the halorhodopsins we have examined to date inactivate by roughly 30% every 15s under typical illumination conditions ($1\text{--}10\text{mW/mm}^2$, of yellow, e.g., 593nm, light; see Fig. 3a and b), consistent with the photocycle topology shown in Fig. 4a (Chow et al., 2010; Han and Boyden, 2007). Further, recovery in the dark of halorhodopsins from this inactivated state is slow, taking tens of minutes for full recovery (Bamberg et al., 1993; Han and Boyden, 2007; Hegemann et al., 1985). Although this rundown may somewhat compromise performance over very long illumination periods, it is possible to drive the molecule out of this inactive state by using UV or blue light (Bamberg et al., 1993; Han and Boyden, 2007; Fig. 3a and b). The entire photocycle is summarized in Fig. 4a, with the limiting time constants noted in the diagram, and with the states denoted by the light absorption properties of the state, as is conventional. The outer circle of the photocycle describes the dominant mode of operation, in which, after photon absorption, a chloride ion is released into the cytoplasm, and then a second




Fig. 3. Kinetic properties of the two major optical neural silencer classes. (a) (i) Halo-mediated hyperpolarizations in a representative current-clamped hippocampal neuron during 15s of continuous yellow light, followed by four 1-s test pulses of yellow light (one every 30s, starting 10s after the end of the first 15-s period of yellow light). (ii) Halo-mediated hyperpolarizations for the same cell exhibited in (i), but when Halo function is facilitated by a 400-ms pulse of blue light in between the 15-s period of yellow light and the first 1-s test pulse. Adapted from Han and Boyden (2007). (b) Population data for blue-light facilitation of Halo recovery. Plotted are the hyperpolarizations elicited by the four 1-s test pulses of yellow light, normalized to the peak hyperpolarization induced by the original 15-s yellow light pulse (mean \pm std. err.). Black dots represent experiments when no blue light pulse was delivered (as in ai). Open blue dots represent experiments when 400ms of blue light was delivered to facilitate recovery (as in aii). Adapted from Han and Boyden (2007). (c) Raw current trace of a neuron lentivirally infected with Arch, illuminated by a 15-s light pulse (575 \pm 25nm, irradiance 7.8mW/mm²), followed by 1s test pulses delivered starting 15, 45, 75, 105, and 135s after the end of the 15s light pulse. Adapted from Chow et al. (2010). (d) Population data of averaged Arch photocurrents sampled at the times indicated by the vertical dotted lines that extend into (c). Adapted from Chow et al. (2010).

chloride ion is taken up from the extracellular space, thus fulfilling the chloride pumping action. In the center of the photocycle diagram, however, is a proton-pumping series of states that involve the long-lasting inactive state described above, here denoted HR410. This HR410 state is the inactivated state that can be reprimed back to the initial state using blue light.

In contrast to the kinetic properties of halorhodopsins, archaeorhodopsins such as Arch and ArchT spontaneously recover in the

dark, even after extensive illumination (Fig. 3c and d). The time constant of recovery is in the range of tens of seconds, in contrast to that of halorhodopsins, which is in the range of tens of minutes. Although archaeorhodopsin photocycles are still relatively uncharacterized compared to those of earlier-discovered opsins, some insight into proton pump operation can be derived from consideration of the *H. salinarum* bacteriorhodopsin photocycle (Fig. 4b). The topology of the photocycle shares some similarities with that of




Fig. 4. Photocycle models for the two major optical neural silencer classes. (a) The halorhodopsin photocycle at high-power continuous illumination on the timescale of a typical photocycle (i.e., conditions used for neural silencing, greater than few milliseconds). The HR410 intermediate is the origin of long-lived inactivation in neural silencing (e.g., Bamberg et al., 1993; Han and Boyden, 2007). (b) The photocycle of the *H. salinarum* bacteriorhodopsin. The photocycle has been simplified to reflect the dominant photocycle at large continuous illumination on the timescale of a typical photocycle (i.e., conditions used for neural silencing, greater than few milliseconds). The M412 alternate intermediate is the origin of long-lived inactivation with bacteriorhodopsin. In contrast, Arch spontaneously quickly recovers from this state in the dark.

halorhodopsins, although many of the timescales are significantly different.

Toward the future

Improvements in optogenetic silencers are rapidly transpiring, driven by the burgeoning amount of microbial genome being sequenced as well as the influx of bioengineering approaches such as mutagenesis and protein engineering. It is likely that the kinetic and amplitude properties of these molecules will continue to improve, given that only a small part of genomic space has been explored, and only a small number of mutations

and protein changes (e.g., the appending of Kir2.1 trafficking sequences) have been explored. From a scientific standpoint, the optical silencers have been broadly applied in the service of assessing the causal role that specific neurons play in the behaviors and neural computations implemented by brain circuits. Now that these molecules have been safely and efficaciously used to modulate neural activity in awake macaque (Diester et al., 2011; Han et al., 2009a), it is also intriguing to ponder whether new neural control prosthetics will arise, capable of silencing overactive neurons or deleting pathological neural activity patterns, in human patients, in the future.

Acknowledgments

E. S. B. acknowledges funding by the NIH Director's New Innovator Award (DP2OD002002) as well as NIH Grants 1R01NS075421, 1R01DA029639, 1RC1MH088182, 1RC2DE020919, 1R01NS067199, and 1R43NS070453; the NSF CAREER award as well as NSF Grants EFRI 0835878, DMS 0848804, and DMS 1042134; Benesse Foundation, Jerry and Marge Burnett, Department of Defense CDMRP Post-Traumatic Stress Disorder Program, Google, Harvard/MIT Joint Grants Program in Basic Neuroscience, Human Frontiers Science Program, MIT Alumni Class Funds, MIT Intelligence Initiative, MIT McGovern Institute and the McGovern Institute Neurotechnology Award Program, MIT Media Lab, MIT Mind-Machine Project, MIT Neurotechnology Fund, NARSAD, Paul Allen Distinguished Investigator Award, Alfred P. Sloan Foundation, SFN Research Award for Innovation in Neuroscience, and the Wallace H. Coulter Foundation.

The authors declare no competing financial interests.

References

- Adamantidis, A. R., Zhang, F., Aravanis, A. M., Deisseroth, K., & de Lecea, L. (2007). Neural substrates of awakening probed with optogenetic control of hypocretin neurons. *Nature*, *450*(7168), 420–424.
- Antón, J., Peña, A., Valens, M., Santos, F., Glöckner, F. O., Bauer, M., et al. (2005). *Salinibacter ruber*: Genomics and biogeography. In N. Gunde-Cimerman, A. Plemenitas & A. Oren (Eds.), *Adaptation to life in high salt concentrations in archaea, bacteria and eukarya* (pp. 257–266). Dordrecht: Kluwer Academic Publishers.
- Aravanis, A. M., Wang, L. P., Zhang, F., Meltzer, L. A., Mogri, M. Z., Schneider, M. B., et al. (2007). An optical neural interface: In vivo control of rodent motor cortex with integrated fiberoptic and optogenetic technology. *Journal of Neural Engineering*, *4*(3), S143–S156.
- Atasoy, D., Aponte, Y., Su, H. H., & Sternson, S. M. (2008). A FLEX switch targets Channelrhodopsin-2 to multiple cell types for imaging and long-range circuit mapping. *Journal of Neuroscience*, *28*(28), 7025–7030.
- Balashov, S. P., Imasheva, E. S., Boichenko, V. A., Anton, J., Wang, J. M., & Lanyi, J. K. (2005). Xanthorhodopsin: A proton pump with a light-harvesting carotenoid antenna. *Science*, *309*(5743), 2061–2064.
- Bamberg, E., Tittor, J., & Oesterhelt, D. (1993). Light-driven proton or chloride pumping by halorhodopsin. *Proceedings of the National Academy of Sciences of the United States of America*, *90*(2), 639–643.
- Beja, O., Aravind, L., Koonin, E. V., Suzuki, M. T., Hadd, A., Nguyen, L. P., et al. (2000). Bacterial rhodopsin: Evidence for a new type of phototrophy in the sea. *Science*, *289*(5486), 1902–1906.
- Beja, O., Spudich, E. N., Spudich, J. L., Leclerc, M., & DeLong, E. F. (2001). Proteorhodopsin phototrophy in the ocean. *Nature*, *411*(6839), 786–789.
- Berthold, P., Tsunoda, S. P., Ernst, O. P., Mages, W., Gradmann, D., & Hegemann, P. (2008). Channelrhodopsin-1 initiates phototaxis and photophobic responses in *Chlamydomonas* by immediate light-induced depolarization. *The Plant Cell*, *20*(6), 1665–1677.
- Bi, A., Cui, J., Ma, Y. P., Olshevskaya, E., Pu, M., Dizhoor, A. M., et al. (2006). Ectopic expression of a microbial-type rhodopsin restores visual responses in mice with photoreceptor degeneration. *Neuron*, *50*(1), 23–33.
- Blanck, A., & Oesterhelt, D. (1987). The halo-opsin gene. II. Sequence, primary structure of halorhodopsin and comparison with bacteriorhodopsin. *EMBO Journal*, *6*(1), 265–273.
- Boyden, E. S. (2011). A history of optogenetics: The development of tools for controlling brain circuits with light. *FI000 Biology Reports*, *3*, 11.
- Boyden, E. S., Zhang, F., Bamberg, E., Nagel, G., & Deisseroth, K. (2005). Millisecond-timescale, genetically targeted optical control of neural activity. *Nature Neuroscience*, *8*(9), 1263–1268.
- Braiman, M. S., Stern, L. J., Chao, B. H., & Khorana, H. G. (1987). Structure-function studies on bacteriorhodopsin. IV. Purification and renaturation of bacterio-opsin polypeptide expressed in *Escherichia coli*. *Journal of Biological Chemistry*, *262*(19), 9271–9276.
- Brown, L. S. (2004). Fungal rhodopsins and opsin-related proteins: Eukaryotic homologues of bacteriorhodopsin with unknown functions. *Photochemical and Photobiological Sciences*, *3*(6), 555–565.
- Brown, L. S., Needleman, R., & Lanyi, J. K. (1996). Interaction of proton and chloride transfer pathways in recombinant bacteriorhodopsin with chloride transport activity: Implications for the chloride translocation mechanism. *Biochemistry*, *35*(50), 16048–16054.
- Campagnola, L., Wang, H., & Zylka, M. J. (2008). Fiber-coupled light-emitting diode for localized photostimulation of neurons expressing channelrhodopsin-2. *Journal of Neuroscience Methods*, *169*(1), 27–33.
- Chan, S. C., Bernstein, J., & Boyden, E. (2010). Scalable fluidic injector arrays for viral targeting of intact 3-D brain circuits. *Journal of Visualized Experiments*, *35*. pii: 1489. doi: 10.3791/1489.

- Chhatwal, J. P., Hammack, S. E., Jasnow, A. M., Rainnie, D. G., & Ressler, K. J. (2007). Identification of cell-type-specific promoters within the brain using lentiviral vectors. *Gene Therapy*, *14*(7), 575–583.
- Chow, B. Y., Chuong, A. S., Klapoetke, N. C., & Boyden, E. S. (2011a). Synthetic physiology strategies for adapting tools from nature for genetically targeted control of fast biological processes. *Methods in Enzymology*, *497*, 425–443.
- Chow, B. Y., Han, X., Bernstein, J. G., Monahan, P. E., & Boyden, E. S. (2011b). Light-activated ion pumps and channels for temporally precise optical control of activity in genetically targeted neurons. In J. J. Chambers & R. H. Kramer (Eds.), *Photosensitive molecules for controlling biological function* (pp. 99–132). New York: Humana Press.
- Chow, B. Y., Han, X., Dobry, A. S., Qian, X., Chuong, A. S., Li, M., et al. (2010). High-performance genetically targetable optical neural silencing by light-driven proton pumps. *Nature*, *463*(7277), 98–102.
- Diester, I., Kaufman, M. T., Mogri, M., Pashaie, R., Goo, W., Yizhar, O., et al. (2011). An optogenetic toolbox designed for primates. *Nature Neuroscience*, *14*(3), 387–397.
- Dittgen, T., Nimmerjahn, A., Komai, S., Licznarski, P., Waters, J., Margrie, T. W., et al. (2004). Lentivirus-based genetic manipulations of cortical neurons and their optical and electrophysiological monitoring in vivo. *Proceedings of the National Academy of Sciences of the United States of America*, *101*(52), 18206–18211.
- Douglass, A. D., Kraves, S., Deisseroth, K., Schier, A. F., & Engert, F. (2008). Escape behavior elicited by single, channelrhodopsin-2-evoked spikes in zebrafish somatosensory neurons. *Current Biology*, *18*(15), 1133–1137.
- Enami, N., Yoshimura, K., Murakami, M., Okumura, H., Ihara, K., & Kouyama, T. (2006). Crystal structures of archaeal rhodopsin-1 and -2: Common structural motif in archaeal light-driven proton pumps. *Journal of Molecular Biology*, *358*(3), 675–685.
- Essen, L. O. (2002). Halorhodopsin: Light-driven ion pumping made simple? *Current Opinion in Structural Biology*, *12*(4), 516–522.
- Farah, N., Reutsky, I., & Shoham, S. (2007). Patterned optical activation of retinal ganglion cells. *Conference Proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society, 2007*, 6369–6371.
- Friedrich, T., Geibel, S., Kalmbach, R., Chizhov, I., Ataka, K., Heberle, J., et al. (2002). Proteorhodopsin is a light-driven proton pump with variable vectoriality. *Journal of Molecular Biology*, *321*(5), 821–838.
- Gilles-Gonzalez, M. A., Engelman, D. M., & Khorana, H. G. (1991). Structure-function studies of bacteriorhodopsin XV. Effects of deletions in loops B-C and E-F on bacteriorhodopsin chromophore and structure. *Journal of Biological Chemistry*, *266*(13), 8545–8550.
- Gradinaru, V., Thompson, K. R., & Deisseroth, K. (2008). eNpHR: A *Natronomonas* halorhodopsin enhanced for optogenetic applications. *Brain Cell Biology*, *36*(1–4), 129–139.
- Gradinaru, V., Thompson, K. R., Zhang, F., Mogri, M., Kay, K., Schneider, M. B., et al. (2007). Targeting and readout strategies for fast optical neural control in vitro and in vivo. *Journal of Neuroscience*, *27*(52), 14231–14238.
- Gradinaru, V., Zhang, F., Ramakrishnan, C., Mattis, J., Prakash, R., Diester, I., et al. (2010). Molecular and cellular approaches for diversifying and extending optogenetics. *Cell*, *141*(1), 154–165.
- Greenberg, K. P., Pham, A., & Werblin, F. S. (2011). Differential targeting of optical neuromodulators to ganglion cell soma and dendrites allows dynamic control of center-surround antagonism. *Neuron*, *69*(4), 713–720.
- Guo, Z. V., Hart, A. C., & Ramanathan, S. (2009). Optical interrogation of neural circuits in *Caenorhabditis elegans*. *Nature Methods*, *6*(12), 891–896.
- Han, X., & Boyden, E. S. (2007). Multiple-color optical activation, silencing, and desynchronization of neural activity, with single-spike temporal resolution. *PLoS One*, *2*(3), e299.
- Han, X., Chow, B. Y., Zhou, H., Klapoetke, N., Chuong, A., Rajimehr, R., et al. (2011). A high-light sensitivity optical neural silencer: Development and application to optogenetic control of non-human primate cortex. *Frontiers in Systems Neuroscience*, *5*, 18.
- Han, X., Qian, X., Bernstein, J. G., Zhou, H. H., Franzesi, G. T., Stern, P., et al. (2009a). Millisecond-timescale optical control of neural dynamics in the nonhuman primate brain. *Neuron*, *62*(2), 191–198.
- Han, X., Qian, X., Stern, P., Chuong, A. S., & Boyden, E. S. (2009b). Informational lesions: Optical perturbation of spike timing and neural synchrony via microbial opsin gene fusions. *Frontiers in Molecular Neuroscience*, doi:10.3389/neuro.02.012.2009.
- Hegemann, P., Oesterhelt, D., & Steiner, M. (1985). The photocycle of the chloride pump halorhodopsin. I: Azide-catalyzed deprotonation of the chromophore is a side reaction of photocycle intermediates inactivating the pump. *EMBO Journal*, *4*(9), 2347–2350.
- Henderson, R., & Schertler, G. F. (1990). The structure of bacteriorhodopsin and its relevance to the visual opsins and other seven-helix G-protein coupled receptors. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *326*(1236), 379–389.
- Huber, D., Petreanu, L., Ghitani, N., Ranade, S., Hromadka, T., Mainen, Z., et al. (2008). Sparse optical microstimulation in barrel cortex drives learned behaviour in freely moving mice. *Nature*, *451*(7174), 61–64.
- Ihara, K., Umemura, T., Katagiri, I., Kitajima-Ihara, T., Sugiyama, Y., Kimura, Y., et al. (1999). Evolution of the archaeal rhodopsins: Evolution rate changes by gene duplication and functional differentiation. *Journal of Molecular Biology*, *285*(1), 163–174.
- Ishizuka, T., Kakuda, M., Araki, R., & Yawo, H. (2006). Kinetic evaluation of photosensitivity in genetically

- engineered neurons expressing green algae light-gated channels. *Neuroscience Research*, 54(2), 85–94.
- Kelemen, B. R., Du, M., & Jensen, R. B. (2003). Proteorhodopsin in living color: Diversity of spectral properties within living bacterial cells. *Biochimica et Biophysica Acta*, 1618(1), 25–32.
- Kim, S. Y., Waschuk, S. A., Brown, L. S., & Jung, K. H. (2008). Screening and characterization of proteorhodopsin color-tuning mutations in *Escherichia coli* with endogenous retinal synthesis. *Biochimica et Biophysica Acta*, 1777(6), 504–513.
- Klare, J. P., Chizhov, I., & Engelhard, M. (2008). Microbial rhodopsins: Scaffolds for ion pumps, channels, and sensors. *Reviews and Problems in Cell Differentiation*, 45, 73–122.
- Kolbe, M., Besir, H., Essen, L. O., & Oesterhelt, D. (2000). Structure of the light-driven chloride pump halorhodopsin at 1.8 Å resolution. *Science*, 288(5470), 1390–1396.
- Kuhlman, S. J., & Huang, Z. J. (2008). High-resolution labeling and functional manipulation of specific neuron types in mouse brain by Cre-activated viral gene expression. *PLoS One*, 3(4), e2005.
- Lagali, P. S., Balya, D., Awatramani, G. B., Munch, T. A., Kim, D. S., Busskamp, V., et al. (2008). Light-activated channels targeted to ON bipolar cells restore visual function in retinal degeneration. *Nature Neuroscience*, 11(6), 667–675.
- Lanyi, J. K. (1986). Halorhodopsin: A light-driven chloride ion pump. *Annual Review of Biophysics and Biophysical Chemistry*, 15(1), 11–28.
- Lanyi, J. K. (2004). Bacteriorhodopsin. *Annual Review of Physiology*, 66(1), 665–688.
- Lanyi, J. K., Duschl, A., Hatfield, G. W., May, K., & Oesterhelt, D. (1990). The primary structure of a halorhodopsin from *Natronobacterium pharaonis*. Structural, functional and evolutionary implications for bacterial rhodopsins and halorhodopsins. *Journal of Biological Chemistry*, 265(3), 1253–1260.
- Li, X., Gutierrez, D. V., Hanson, M. G., Han, J., Mark, M. D., Chiel, H., et al. (2005). Fast noninvasive activation and inhibition of neural and network activity by vertebrate rhodopsin and green algae channelrhodopsin. *Proceedings of the National Academy of Sciences of the United States of America*, 102(49), 17816–17821.
- Lin, J. Y., Lin, M. Z., Steinbach, P., & Tsien, R. Y. (2009). Characterization of engineered channelrhodopsin variants with improved properties and kinetics. *Biophysical Journal*, 96(5), 1803–1814.
- Luecke, H., Schobert, B., Richter, H. T., Cartailler, J. P., & Lanyi, J. K. (1999). Structure of bacteriorhodopsin at 1.55 Å resolution. *Journal of Molecular Biology*, 291(4), 899–911.
- Lukashev, E. P., Govindjee, R., Kono, M., Ebrey, T. G., Sugiyama, Y., & Mukohata, Y. (1994). pH dependence of the absorption spectra and photochemical transformations of the archaeorhodopsins. *Photochemistry and Photobiology*, 60(1), 69–75.
- Marinetti, T., Subramaniam, S., Mogi, T., Marti, T., & Khorana, H. G. (1989). Replacement of aspartic residues 85, 96, 115, or 212 affects the quantum yield and kinetics of proton release and uptake by bacteriorhodopsin. *Proceedings of the National Academy of Sciences of the United States of America*, 86(2), 529–533.
- Marti, T., Otto, H., Mogi, T., Rosselet, S. J., Heyn, M. P., & Khorana, H. G. (1991). Bacteriorhodopsin mutants containing single substitutions of serine or threonine residues are all active in proton translocation. *Journal of Biological Chemistry*, 266(11), 6919–6927.
- Ming, M., Lu, M., Balashov, S. P., Ebrey, T. G., Li, Q., & Ding, J. (2006). pH dependence of light-driven proton pumping by an archaeorhodopsin from Tibet: Comparison with bacteriorhodopsin. *Biophysical Journal*, 90(9), 3322–3332.
- Mogi, T., Marti, T., & Khorana, H. G. (1989a). Structure-function studies on bacteriorhodopsin. IX. Substitutions of tryptophan residues affect protein-retinal interactions in bacteriorhodopsin. *Journal of Biological Chemistry*, 264(24), 14197–14201.
- Mogi, T., Stern, L. J., Chao, B. H., & Khorana, H. G. (1989b). Structure-function studies on bacteriorhodopsin. VIII. Substitutions of the membrane-embedded prolines 50, 91, and 186: The effects are determined by the substituting amino acids. *Journal of Biological Chemistry*, 264(24), 14192–14196.
- Mogi, T., Stern, L. J., Hackett, N. R., & Khorana, H. G. (1987). Bacteriorhodopsin mutants containing single tyrosine to phenylalanine substitutions are all active in proton translocation. *Proceedings of the National Academy of Sciences of the United States of America*, 84(16), 5595–5599.
- Mogi, T., Stern, L. J., Marti, T., Chao, B. H., & Khorana, H. G. (1988). Aspartic acid substitutions affect proton translocation by bacteriorhodopsin. *Proceedings of the National Academy of Sciences of the United States of America*, 85(12), 4148–4152.
- Mukohata, Y., Ihara, K., Tamura, T., & Sugiyama, Y. (1999). Halobacterial rhodopsins. *Journal of Biochemistry*, 125(4), 649–657.
- Nagel, G., Brauner, M., Liewald, J. F., Adeishvili, N., Bamberg, E., & Gottschalk, A. (2005). Light activation of channelrhodopsin-2 in excitable cells of *Caenorhabditis elegans* triggers rapid behavioral responses. *Current Biology*, 15(24), 2279–2284.
- Nagel, G., Szellas, T., Huhn, W., Kateriya, S., Adeishvili, N., Berthold, P., et al. (2003). Channelrhodopsin-2, a directly light-gated cation-selective membrane channel. *Proceedings of the National Academy of Sciences of the United States of America*, 100(24), 13940–13945. (PMCID: 283525).
- Okuno, D., Asaumi, M., & Muneyuki, E. (1999). Chloride concentration dependency of the electrogenic activity of halorhodopsin. *Biochemistry*, 38(17), 5422–5429.
- Petreanu, L., Huber, D., Sobczyk, A., & Svoboda, K. (2007). Channelrhodopsin-2-assisted circuit mapping of long-range callosal projections. *Nature Neuroscience*, 10(5), 663–668.
- Petreanu, L., Mao, T., Sternson, S. M., & Svoboda, K. (2009). The subcellular organization of neocortical excitatory connections. *Nature*, 457(7233), 1142–1145.

- Rickgauer, J. P., & Tank, D. W. (Eds.), (2008). Optimizing two-photon activation of channelrhodopsin-2 for stimulation at cellular resolution. *Neuroscience*. Washington, DC: Society for Neuroscience.
- Rudiger, M., & Oesterhelt, D. (1997). Specific arginine and threonine residues control anion binding and transport in the light-driven chloride pump halorhodopsin. *EMBO Journal*, *16*(13), 3813–3821.
- Schroll, C., Riemensperger, T., Bucher, D., Ehmer, J., Voller, T., Erbguth, K., et al. (2006). Light-induced activation of distinct modulatory neurons triggers appetitive or aversive learning in *Drosophila* larvae. *Current Biology*, *16* (17), 1741–1747.
- Seki, A., Miyauchi, S., Hayashi, S., Kikukawa, T., Kubo, M., Demura, M., et al. (2007). Heterologous expression of Pharaonis halorhodopsin in *Xenopus laevis* oocytes and electrophysiological characterization of its light-driven Cl⁻ pump activity. *Biophysical Journal*, *92*(7), 2559–2569.
- Subramaniam, S., Greenhalgh, D. A., & Khorana, H. G. (1992). Aspartic acid 85 in bacteriorhodopsin functions both as proton acceptor and negative counterion to the Schiff base. *Journal of Biological Chemistry*, *267*(36), 25730–25733.
- Tan, W., Janczewski, W. A., Yang, P., Shao, X. M., Callaway, E. M., & Feldman, J. L. (2008). Silencing pre-Botzinger complex somatostatin-expressing neurons induces persistent apnea in awake rat. *Nature Neuroscience*, *11*(5), 538–540.
- Tittor, J., Haupts, U., Haupts, C., Oesterhelt, D., Becker, A., & Bamberg, E. (1997). Chloride and proton transport in bacteriorhodopsin mutant D85T: Different modes of ion translocation in a retinal protein. *Journal of Molecular Biology*, *271*(3), 405–416.
- Tittor, J., Oesterhelt, D., & Bamberg, E. (1995). Bacteriorhodopsin mutants D85N, D85T and D85, 96N as proton pumps. *Biophysical Chemistry*, *56*(1–2), 153–157.
- Toni, N., Laplagne, D. A., Zhao, C., Lombardi, G., Ribak, C. E., Gage, F. H., et al. (2008). Neurons born in the adult dentate gyrus form functional synapses with target cells. *Nature Neuroscience*, *11*(8), 901–907.
- Tsunematsu, T., Kilduff, T. S., Boyden, E. S., Takahashi, S., Tominaga, M., & Yamanaka, A. (2011). Acute optogenetic silencing of orexin/hypocretin neurons induces slow-wave sleep in mice. *Journal of Neuroscience*, *31*(29), 10529–10539.
- Tsunoda, S. P., Ewers, D., Gazzarrini, S., Moroni, A., Gradmann, D., & Hegemann, P. (2006). H⁺-pumping rhodopsin from the marine alga *acetabularia*. *Biophysical Journal*, *91*(4), 1471–1479.
- Varo, G., Brown, L. S., Sasaki, J., Kandori, H., Maeda, A., Needleman, R., et al. (1995a). Light-driven chloride ion transport by halorhodopsin from *Natronobacterium pharaonis*. 1. The photochemical cycle. *Biochemistry*, *34* (44), 14490–14499.
- Varo, G., Needleman, R., & Lanyi, J. K. (1995b). Light-driven chloride ion transport by halorhodopsin from *Natronobacterium pharaonis*. 2. Chloride release and uptake, protein conformation change, and thermodynamics. *Biochemistry*, *34*(44), 14500–14507.
- Wang, H., Peca, J., Matsuzaki, M., Matsuzaki, K., Noguchi, J., Qiu, L., et al. (2007). High-speed mapping of synaptic connectivity using photostimulation in Channelrhodopsin-2 transgenic mice. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(19), 8143–8148.
- Waschuk, S. A., Bezerra, A. G., Shi, L., & Brown, L. S. (2005). Leptosphaeria rhodopsin: Bacteriorhodopsin-like proton pump from a eukaryote. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(19), 6879–6883.
- Wentz, C. T., Bernstein, J. G., Monahan, P., Guerra, A., Rodriguez, A., & Boyden, E. S. (2011). A wirelessly powered and controlled device for optical neural control of freely-behaving animals. *Journal of Neural Engineering*, *8* (4), 046021.
- Yoshimura, K., & Kouyama, T. (2008). Structural role of bacterioruberin in the trimeric structure of archaeorhodopsin-2. *Journal of Molecular Biology*, *375*(5), 1267–1281.
- Zhang, F., Wang, L. P., Boyden, E. S., & Deisseroth, K. (2006). Channelrhodopsin-2 and optical control of excitable cells. *Nature Methods*, *3*(10), 785–792.
- Zhang, F., Wang, L. P., Brauner, M., Liewald, J. F., Kay, K., Watzke, N., et al. (2007). Multimodal fast optical interrogation of neural circuitry. *Nature*, *446*(7136), 633–639.
- Zhao, S., Cunha, C., Zhang, F., Liu, Q., Gloss, B., Deisseroth, K., et al. (2008). Improved expression of halorhodopsin for light-induced silencing of neuronal activity. *Brain Cell Biology*, *36*(1–4), 141–154.
- Zorzos, A. N., Boyden, E. S., & Fonstad, C. G. (2010). Multi-wavelength implantable probe for light delivery to sets of distributed brain targets. *Optics Letters*, *35*(24), 4133–4135.